



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Porphyromonas gingivalis: an invasive and evasive opportunistic oral pathogen

Bostanci, Nagihan ; Belibasakis, Georgios N

Abstract: *Porphyromonas gingivalis* is a Gram-negative oral anaerobe that is involved in the pathogenesis of periodontitis, an inflammatory disease that destroys the tissues supporting the tooth, eventually leading to tooth loss. *Porphyromonas gingivalis* has can locally invade periodontal tissues and evade the host defence mechanisms. In doing so, it utilizes a panel of virulence factors that cause deregulation of the innate immune and inflammatory responses. The present review discusses the invasive and evasive strategies of *P. gingivalis* and the role of its major virulence factors in these, namely lipopolysaccharide, capsule, gingipains and fimbriae. Moreover, the role of *P. gingivalis* as a 'keystone' biofilm species in orchestrating a host response, is highlighted.

DOI: <https://doi.org/10.1111/j.1574-6968.2012.02579.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-63941>

Journal Article

Published Version

Originally published at:

Bostanci, Nagihan; Belibasakis, Georgios N (2012). *Porphyromonas gingivalis*: an invasive and evasive opportunistic oral pathogen. *FEMS Microbiology Letters*, 333(1):1-9.

DOI: <https://doi.org/10.1111/j.1574-6968.2012.02579.x>

MINIREVIEW

***Porphyromonas gingivalis*: an invasive and evasive opportunistic oral pathogen**

Nagihan Bostanci¹ & Georgios N. Belibasakis²

¹Oral Translational Research, Institute of Oral Biology, Center of Dental Medicine, Faculty of Medicine, University of Zürich, Zürich, Switzerland; and ²Oral Microbiology and Immunology, Institute of Oral Biology, Center of Dental Medicine, Faculty of Medicine, University of Zürich, Zürich, Switzerland

Correspondence: Georgios N. Belibasakis, Oral Microbiology and Immunology, Institute of Oral Biology, Center of Dental Medicine, Faculty of Medicine, University of Zürich, Plattenstrasse 11, 8032 Zürich, Switzerland. Tel.: +41 44 634 3306; fax: +41 44 634 4310; e-mail: george.belibasakis@zzm.uzh.ch

Received 25 January 2012; revised 18 April 2012; accepted 20 April 2012.
Final version published online 28 May 2012.

DOI: 10.1111/j.1574-6968.2012.02579.x

Editor: Michael Mourez

Keywords

Porphyromonas gingivalis; inflammation; immune response; periodontitis.

Abstract

Porphyromonas gingivalis is a Gram-negative oral anaerobe that is involved in the pathogenesis of periodontitis, an inflammatory disease that destroys the tissues supporting the tooth, eventually leading to tooth loss. *Porphyromonas gingivalis* can locally invade periodontal tissues and evade the host defence mechanisms. In doing so, it utilizes a panel of virulence factors that cause deregulation of the innate immune and inflammatory responses. The present review discusses the invasive and evasive strategies of *P. gingivalis* and the role of its major virulence factors in these, namely lipopolysaccharide, capsule, gingipains and fimbriae. Moreover, the role of *P. gingivalis* as a 'keystone' biofilm species in orchestrating a host response, is highlighted.

Introduction

***Porphyromonas gingivalis* and association with periodontal disease**

Periodontal disease, or periodontitis, is defined as a bacterially induced inflammatory disease of the tooth-supporting (periodontal) tissues. Although more than 700 bacterial species can colonize the oral cavity (Aas *et al.*, 2005), only a handful of those are highly implicated in the disease (Paster *et al.*, 2006). *Porphyromonas gingivalis* is the species most highly associated with the chronic form of periodontitis, and can be detected in up to 85% of the disease sites (Yang *et al.*, 2004). It is detected rarely or at low in numbers in healthy sites. The presence of *P. gingivalis* in a periodontal pocket may predict imminent disease progression (van Winkelhoff *et al.*, 2002) and a significant positive correlation is found between *P. gingivalis* numbers and pocket depth (Kawada *et al.*, 2004). Following periodontal treatment, a reduction of *P. gingivalis* numbers is associated with resolution of

disease at the affected site (Haffajee *et al.*, 1997; Fujise *et al.*, 2002). Moreover, experimental implantation of *P. gingivalis* in animal models induces an inflammatory response and periodontal bone loss (Evans *et al.*, 1992; Hajishengallis *et al.*, 2011). This species possesses a number of potential virulence factors, such as cysteine proteinases (gingipains), lipopolysaccharide (LPS), capsule and fimbriae (Lamont & Jenkinson, 1998). Collectively, due to these properties *P. gingivalis* is considered an 'opportunistic pathogen', in line with the modified Koch's postulates for oral infections, such as periodontal diseases (Socransky, 1979).

Structural and growth characteristics of *P. gingivalis*

Porphyromonas gingivalis is a black-pigmented, assaccharolytic, non-motile Gram-negative species that requires anaerobic conditions for growth, and the presence of heme or hemin and vitamin K in its nutrient milieu. It gains its metabolic energy by fermenting amino acids, a

property decisive for its survival in deep periodontal pockets, where sugars are extremely scarce. When considering its location in multispecies subgingival biofilm communities, *P. gingivalis* is a late colonizer, and hence is found in close proximity to and interacts with the juxtaposing gingival tissue (Kolenbrander *et al.*, 2011; Zijnga *et al.*, 2011). The black pigmentation of *P. gingivalis* colonies observed in blood agar culture is itself associated with the aggregation of heme on its cell surface (Liu *et al.*, 2004; Smalley *et al.*, 2006). This property is somehow connected to its capacity to act as an opportunistic pathogen, as when grown in a heme-limited medium it becomes less virulent (McKee *et al.*, 1986).

Invasion of the host by *P. gingivalis*

As part of its strategies for survival into the host, *P. gingivalis* is able to invade cells and tissues (Yilmaz, 2008), thus avoiding the immune surveillance. *Porphyromonas gingivalis* can actively invade gingival epithelial cells, where it can maintain viability and replicate (Belton *et al.*, 1999; Tribble *et al.*, 2006). This invasive property is dependent on its major fimbriae, which bind to $\beta 1$ integrin on the surface of host cells, an event that causes rearrangements of the actin cytoskeleton to allow internalization (Yilmaz *et al.*, 2002, 2003). *Porphyromonas gingivalis* can also invade macrophages, but within these cells its replication is less active (Wang *et al.*, 2007). This is potentially a strategy for limited exposure to the extracellular environment and evasion of the immune surveillance. Interestingly, once *P. gingivalis* has invaded intracellularly, there are no signs of apoptosis or necrosis (Nakhjiri *et al.*, 2001). It can then actively secrete an ATP-hydrolysing enzyme, thus suppressing ATP-dependent apoptosis (Yilmaz *et al.*, 2008) and allowing its survival in host cells. Subsequently, it can disseminate from cell to cell, through actin cytoskeleton bridges without causing cell death, and spread while avoiding immune surveillance (Yilmaz *et al.*, 2006). Once *P. gingivalis* is established in the cell, it affects cell-cycle pathways and thus accelerates proliferation of gingival epithelial cells, in a fimbriae-dependent fashion (Kuboniwa *et al.*, 2008). This could well constitute a mechanism of expansion of the periodontal pocket epithelium, which is a histopathological feature of periodontitis.

Survival strategies of *P. gingivalis*

It is now well established that *P. gingivalis* is not an aggressor of the inflammatory response, but rather an opportunist that can cross-talk with the host and subvert its defence mechanisms. Using this strategy, *P. gingivalis* prolongs its survival and becomes established in the

periodontal pocket (Hajishengallis *et al.*, 2011). It preferentially deregulates innate immunity, which may in turn disable adaptive immunity (Hajishengallis, 2009; Pathirana *et al.*, 2010). Important representative examples of these abilities are its capacity to degrade human defensins (Carlisle *et al.*, 2009), its resistance to oxidative burst-killing by polymorphonuclear neutrophils (PMNs) (Mydel *et al.*, 2006) and its ability to inhibit 'at will' the production of crucial proinflammatory cytokines (Bostanci *et al.*, 2007a, b). Although *P. gingivalis* has the capacity to stimulate interleukin (IL)-8 production by epithelial cells (Sandros *et al.*, 2000; Asai *et al.*, 2001; Kusumoto *et al.*, 2004), it can also inhibit IL-8 production, resulting in hindered PMN chemotaxis, a phenomenon known as 'chemokine paralysis' (Darveau *et al.*, 1998). *Porphyromonas gingivalis* thereby incapacitates the first line of defence in the periodontal tissues. Moreover, by inhibiting IL-12 production by macrophages, it prevents cytotoxic T-cell activation and therefore bacterial clearance (Hajishengallis *et al.*, 2007). Accordingly, by inhibiting interferon (IFN)- γ production by T cells, it inhibits macrophage bacteriocidal activity and hence bacterial clearance (Pulendran *et al.*, 2001; Hajishengallis *et al.*, 2007). A special relationship is also revealed between *P. gingivalis* and the complement system, as it can suppress its activation, that is by degradation of C3 and capturing of C4b-binding protein, but also by synergizing with C5a via exploiting toll-like receptor (TLR)-2 signalling (Wang *et al.*, 2010). A further interesting point is that whole viable *P. gingivalis* is differentially sensed by the host, compared with its released virulence factors, with the potential to activate distinctive intracellular pathways (Pathirana *et al.*, 2010), or differential cytokine production (Zhou *et al.*, 2005).

Virulence factors of *P. gingivalis*

As an opportunistic pathogen, it is not surprising that *P. gingivalis* possesses a number of virulence factors. These are molecules that can elicit deleterious effects on host cells, essentially the survival 'weapons' of *P. gingivalis*. The main virulence factors discussed here are LPS, capsular polysaccharide (CPS), fimbriae and gingipains.

The LPS of *P. gingivalis*

Like all Gram-negative bacterial species, *P. gingivalis* is sheathed by an LPS, which is an outer membrane component recognized by the host that can trigger intracellular signalling events. The affinity of LPS to its pattern recognition receptors, such as the TLRs and CD14, enables discrimination between commensal and pathogenic species. The *P. gingivalis* LPS is a stimulator of proinflammatory

responses and bone resorption, as demonstrated in experimental animal models (Chiang *et al.*, 1999; Nishida *et al.*, 2001). *In vitro*, it stimulates proinflammatory cytokine production of, for example, IL-1 α , IL-1 β , IL-6, IL-8, IL-18 and tumour necrosis factor (TNF)- α in monocytes (Zhou *et al.*, 2005; Bostanci *et al.*, 2007a, b; Hamed *et al.*, 2009). Yet, *P. gingivalis* LPS exhibits controversial features with regard to the induction of an inflammatory response. Apart from being a weaker cytokine stimulator compared with the LPS of other Gram-negative (i.e. enteropathogenic) species (Liu *et al.*, 2008), it can also antagonize the cytokine-stimulating capacity of other putative pathogens (Bostanci *et al.*, 2007a, b).

Structurally, *P. gingivalis* LPS exhibits unique features compared with the LPS of other species. These include differences in the structure of the O-antigen between *P. gingivalis* strains that can confer antigenic differences (Paramonov *et al.*, 2001, 2009), as well as in the acylation patterns and receptor-activating capacities of the lipid A component. While the lipid A of most Gram-negative species is a strong activator of TLR4 responses, *P. gingivalis* lipid A is predominantly a TLR2 activator and may even act as antagonist to TLR4 (Darveau *et al.*, 2004), dampening the immune responses (Hajishengallis, 2009). When considering further the heterogeneous acylation patterns of *P. gingivalis* lipid A, two forms are predominant: the tetra-acylated and penta-acylated forms. These two structures induce opposing host responses. The penta-acylated lipid A activates TLR4, whereas tetra-acylated lipid A acts as a TLR4 antagonist (Darveau *et al.*, 2004; Nemoto *et al.*, 2006). These changes of *P. gingivalis* lipid A acylation are dependent on microenvironmental conditions. In particular, when hemin availability is high (a condition that reflects inflammation), penta-acylated lipid A is converted into tetra-acylated lipid A (Al-Qutub *et al.*, 2006). Hence, by modifying its lipid A structure according to the microenvironment, *P. gingivalis* may modulate the binding affinity of its LPS to its cognate TLR receptors, subsequently selecting how to affect downstream host immune signalling. Interestingly, a second type of LPS has also been identified in *P. gingivalis*, containing a distinct anionic polysaccharide linked to lipid A, known as A-LPS (Paramonov *et al.*, 2005). A-LPS is required for cell integrity and serum resistance (Shoji *et al.*, 2002; Paramonov *et al.*, 2005; Slaney *et al.*, 2006) and is structurally associated with the Arg-X gingipain (Curtis *et al.*, 1999; Paramonov *et al.*, 2005). It is also a weaker inducer of cytokine responses by human monocytes, as compared with the conventional LPS (Rangarajan *et al.*, 2008). Collectively, the modifications and heterogeneity of *P. gingivalis* LPS can result in opposing actions and immunological deregulation. Strategically, this is in line with the manipulation of host innate immune

responses by this species, to facilitate its adaptation and survival into the host.

The CPS of *P. gingivalis*

A major virulence factor of *P. gingivalis* is considered to be its capsule, also known as CPS or K-antigen (Schifferle *et al.*, 1989; Holt *et al.*, 1999; Farquharson *et al.*, 2000; Aduse-Opoku *et al.*, 2006; Brunner *et al.*, 2010a, b). Based on the capacity of CPS to generate systemic IgG antibody responses, at least six different serotypes have been identified (Laine *et al.*, 1997; Sims *et al.*, 2001). Encapsulated *P. gingivalis* strains are shown to be highly invasive, causing spreading infection in a murine lesion model, whereas nonencapsulated strains induced only localized abscesses (Laine & van Winkelhoff, 1998). Interestingly, immunization with *P. gingivalis* CPS induced a high IgG systemic response (Choi *et al.*, 1998) and reduced *P. gingivalis*-induced alveolar bone loss (Gonzalez *et al.*, 2003). Encapsulated strains of *P. gingivalis* are more resistant to phagocytosis by polymorphonuclear leukocytes than nonencapsulated strains (Sundqvist *et al.*, 1991) and have differential capacities to adhere to gingival epithelial cells (Dierickx *et al.*, 2003). Moreover, differences in CPS serotypes can reflect differential capacities in chemokine stimulation by macrophages (d'Empaire *et al.*, 2006) or cytokine stimulation by dendritic cells (Vernal *et al.*, 2009). Interestingly, a nonencapsulated *P. gingivalis* knockout mutant strain was found to be a more potent inducer of cytokine synthesis by human gingival fibroblasts, as compared with the corresponding wild-type strain, implying a role of CPS in downplaying the innate immune responses (Brunner *et al.*, 2010a, b). Although it is evident that the presence of CPS, or its individual serotypes, could be determinants of the virulence of *P. gingivalis*, the potential involvement of this antigen in the overall deregulation of host responses awaits further clarification.

The fimbriae of *P. gingivalis*

The fimbriae of *P. gingivalis* are thin, filamentous cell-surface protrusions that facilitate its adherence to salivary proteins, extracellular matrix, eukaryotic cells and bacteria of either the same or other species. Through its fimbriae, *P. gingivalis* can thus attach to early colonizing bacteria, and participate in the developing biofilm structure. Type I (major) fimbriae have important roles in colonization and invasion, whereas type II (minor) fimbriae possess a higher proinflammatory capacity (Lamont & Jenkinson, 1998; Amano *et al.*, 2004; Hajishengallis *et al.*, 2008). Interestingly, however, *P. gingivalis* strains W50 and W83 that lack major fimbriae are still invasive, as

demonstrated in experimental subcutaneous abscess models (Inaba *et al.*, 2008). A particular role of fimbriae is revealed in the induction of bone destruction in experimental periodontitis models. To this extent, infection of rats with nonfimbriated *P. gingivalis* strains exhibited reduced periodontal bone loss, compared with infection with fimbriated strains (Jotwani & Cutler, 2004). Moreover, immunization against *P. gingivalis* fimbriae protected against bone loss in gnotobiotic rats (Malek *et al.*, 1994; Sharma *et al.*, 2001). Other properties of both major and minor fimbriae are the induction of proinflammatory cytokines and production of matrix metalloproteinases (MMPs), such as IL-1, IL-6, IL-8, TNF- α and MMP-9, by various host cells (Jotwani *et al.*, 2010; Ogawa *et al.*, 1994; Pollreis *et al.*, 2010; Takahashi *et al.*, 2006).

Porphyromonas gingivalis fimbriae can signal through either TLR2 or TLR4. Activation of TLR2 by fimbriae results in a differential signalling pattern compared with activation by *P. gingivalis* LPS (Hajishengallis *et al.*, 2006). Fimbriae can directly induce two distinct signalling pathways, one that mediates production of proinflammatory cytokines, such as IL-6 and TNF- α , and another that mediates the expression of cell adhesion molecules, such as ICAM-1 (Hajishengallis *et al.*, 2009). On the other hand, signalling through TLR4 requires an additional costimulation of CD14 and MD-2 (Davey *et al.*, 2008). Interestingly, major fimbriae can exploit TLR2 signalling in order to interact with complement receptor 3 (CR3), in a novel 'inside-out' signalling pattern (Hajishengallis *et al.*, 2007; Wang *et al.*, 2007). This interaction activates the binding capacity of CR3 and allows for internalization of *P. gingivalis* in macrophages and reduction of IL-12 production, which may collectively inhibit bacterial clearance (Hajishengallis *et al.*, 2007).

The gingipains of *P. gingivalis*

Gingipains are a group of cell surface cysteine proteinases of *P. gingivalis* that can also be present in secreted soluble form. They account for 85% of the total proteolytic activity of *P. gingivalis* (Potempa *et al.*, 1997). Based on their substrate specificity, they are divided into arginine-specific (Arg-X) and lysine-specific (Lys-X) gingipains (Curtis *et al.*, 2001; Guo *et al.*, 2010). Arg-X gingipains have trypsin-like activity, and can degrade extracellular matrix components, including the integrin–fibronectin-binding, cytokine, immunoglobulin and complement factors. There are two types of Arg-X gingipains, namely RgpA, which contains a proteolytic and an adhesion domain, and RgpB, which contains only the proteolytic domain. There is one type of Lys-X gingipain, Kgp, which contains both a proteolytic and an adhesion domain. There are sequence similarities between the adhesion domains of Kgp and RgpA (Curtis *et al.*, 2001).

The gingipains have multiple effects on the molecular components of the immune response, and as such they can deregulate these responses. For instance, they can cleave several T-cell receptors, such as CD2, CD4 and CD8 (Kitamura *et al.*, 2002), thereby hampering the cell-mediated immune response. They can also stimulate expression of protease-activated receptors in neutrophils (Lourbakos *et al.*, 1998), gingival epithelial cells (Lourbakos *et al.*, 2001), gingival fibroblasts and T cells (Belibasakis *et al.*, 2010), which are crucial for the induction of cytokine responses and the establishment of chronic inflammation in periodontitis (Holzhausen *et al.*, 2010; Fagundes *et al.*, 2011). Gingipains can also stimulate IL-6 production by oral epithelial cells (Lourbakos *et al.*, 2001) and IL-8 production by gingival fibroblasts (Oido-Mori *et al.*, 2001), enhancing the inflammatory responses. However, they can also proteolytically inactivate both anti-inflammatory (IL-4, IL-5) and pro-inflammatory (IL-12, IFN- γ) cytokines (Yun *et al.*, 1999, 2001, 2002; Tam *et al.*, 2009).

A number of particularly interesting effects are exerted by the gingipains on components of the complement system. Arg-X gingipains can cleave the C5 molecule, resulting in release of its C5a component, which is crucial for enhancing the recruitment of PMNs (Wingrove *et al.*, 1992; Imamura *et al.*, 2001). On the other hand, Lys-X can inactivate the C5a receptor on PMNs, an action that may actually impair their recruitment (Jagels *et al.*, 1996a, b). Along this line, the Arg-X gingipains can degrade the C3 molecule, potentially contributing to decreased bacterial opsonization (Schenkein *et al.*, 1995). This property could confer increased resistance of *P. gingivalis* to bactericidal activity.

Apart from their effect on immune responses, gingipains may also be involved in the binding of *P. gingivalis* to host cells, as Rgp–Kgp complexes have been shown to mediate adherence on gingival epithelial cells and gingival fibroblasts (Chen *et al.*, 2001; Grenier *et al.*, 2003; Andrian *et al.*, 2004). Interestingly, when *P. gingivalis* intracellularly invades gingival epithelial cells, expression of gingipain is downregulated (Xia *et al.*, 2007).

Gingipains may also affect vascular permeability and bleeding at the periodontal site. They can proteolytically activate plasma kallikrein and bradykinin, or alternatively increase the release of thrombin and prothrombin, which can result in increased vascular permeability and PMN influx (Imamura *et al.*, 1994, 1995a). Moreover, by degrading fibrinogen (Scott *et al.*, 1993), they may contribute to inhibition of blood coagulation and increase bleeding at the site (Imamura *et al.*, 1995b), thus enhancing the availability of hemin required for *P. gingivalis* growth.

Collectively, studies in various experimental systems indicate that gingipains have seemingly contradicting

actions on the innate immune responses, hampering interpretation of their role in the pathogenesis of periodontitis. Nevertheless, such differences may be reconciled by the existence of a concentration gradient of gingipains in the tissue (Pathirana *et al.*, 2010). Closer to the gingival epithelial barrier where the biofilm resides, gingipain concentrations are high, causing degradation or deregulation of various components of the immune response. This hampers bacterial clearance and facilitates bacterial invasion. Nevertheless, deeper into the gingival connective tissue, gingipain concentrations become gradually lower and stimulate, rather than inhibit, inflammation. This may in turn induce connective tissue and bone destruction, which are the hallmarks of periodontitis.

Conclusions

It is evident that *P. gingivalis* has developed mechanisms to invade and persist into the host, by astutely adapting to its local niche. Its paradoxically opposing (stimulatory vs. inhibiting) effects on innate immune and inflammatory responses aim to subvert host defence mechanisms, in order to facilitate its survival in the tissues (Hajishengallis, 2009; Hajishengallis & Lambris, 2011). The net effect of this deregulated equilibrium is likely to determine if site-specific disease progresses beyond or remains at stationary phase. Whether inflammation is beneficial for *P. gingivalis* may depend on the stage of its establishment in the host (Hajishengallis, 2009; Pathirana *et al.*, 2010). At early stages, suppression of inflammation and evasion of host recognition would aid *P. gingivalis* in colonizing, invading and establishing at the targeted site. At later stages, once *P. gingivalis* is well established, inducing inflammation may facilitate its increased demands in nutrients. Alternatively, *P. gingivalis* may induce a 'non-productive inflammation', one that fails to eliminate it, yet is sufficient to induce mediators of tissue destruction (Hajishengallis, 2009).

Finally, as periodontitis is of polymicrobial nature, it is reasonable to consider the role of different bacterial species within the context of (subgingival) biofilm communities. Hence, *P. gingivalis* is likely to function in concerted action with other species, to their mutual benefit. For instance, complement manipulation by *P. gingivalis* may denote a coevolution strategy to support other species present in the biofilm, which may reciprocally provide further colonization opportunities and nutrient availability to *P. gingivalis*. Subsequent changes in the local microenvironment can differentially regulate expression of its virulence factors, and hence the proinflammatory or anti-inflammatory potentials of *P. gingivalis*. This is strongly indicated by recent evidence demonstrating that even at low abundance, this species qualitatively and

quantitatively affects the composition of the oral commensal microbiota, which are in turn required for *P. gingivalis*-induced inflammatory bone loss (Hajishengallis *et al.*, 2011). For these reasons, *P. gingivalis* is now considered a 'keystone' species in subgingival biofilms (Honda, 2011).

Acknowledgements

This work was supported by the Institute of Oral Biology, Center of Dental Medicine, University of Zürich.

References

- Aas JA, Paster BJ, Stokes LN, Olsen I & Dewhirst FE (2005) Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol* **43**: 5721–5732.
- Aduse-Opoku J, Slaney JM, Hashim A, Gallagher A, Gallagher RP, Rangarajan M, Boutaga K, Laine ML, Van Winkelhoff AJ & Curtis MA (2006) Identification and characterization of the capsular polysaccharide (K-antigen) locus of *Porphyromonas gingivalis*. *Infect Immun* **74**: 449–460.
- Al-Qutub MN, Braham PH, Karimi-Naser LM, Liu X, Genco CA & Darveau RP (2006) Hemin-dependent modulation of the lipid A structure of *Porphyromonas gingivalis* lipopolysaccharide. *Infect Immun* **74**: 4474–4485.
- Amano A, Nakagawa I, Okahashi N & Hamada N (2004) Variations of *Porphyromonas gingivalis* fimbriae in relation to microbial pathogenesis. *J Periodontol Res* **39**: 136–142.
- Andrian E, Grenier D & Rouabhia M (2004) *In vitro* models of tissue penetration and destruction by *Porphyromonas gingivalis*. *Infect Immun* **72**: 4689–4698.
- Asai Y, Ohyama Y, Gen K & Ogawa T (2001) Bacterial fimbriae and their peptides activate human gingival epithelial cells through Toll-like receptor 2. *Infect Immun* **69**: 7387–7395.
- Belibasakis GN, Bostanci N & Reddi D (2010) Regulation of protease-activated receptor-2 expression in gingival fibroblasts and Jurkat T cells by *Porphyromonas gingivalis*. *Cell Biol Int* **34**: 287–292.
- Belton CM, Izutsu KT, Goodwin PC, Park Y & Lamont RJ (1999) Fluorescence image analysis of the association between *Porphyromonas gingivalis* and gingival epithelial cells. *Cell Microbiol* **1**: 215–223.
- Bostanci N, Allaker RP, Belibasakis GN, Rangarajan M, Curtis MA, Hughes FJ & McKay IJ (2007a) *Porphyromonas gingivalis* antagonises *Campylobacter rectus* induced cytokine production by human monocytes. *Cytokine* **39**: 147–156.
- Bostanci N, Allaker R, Johansson U, Rangarajan M, Curtis MA, Hughes FJ & McKay IJ (2007b) Interleukin-1 α stimulation in monocytes by periodontal bacteria: antagonistic effects of *Porphyromonas gingivalis*. *Oral Microbiol Immunol* **22**: 52–60.
- Brunner J, Scheres N, El Idrissi NB, Deng DM, Laine ML, van Winkelhoff AJ & Crielaard W (2010a) The capsule of

- Porphyromonas gingivalis* reduces the immune response of human gingival fibroblasts. *BMC Microbiol* **10**: 5.
- Brunner J, Wittink FR, Jonker MJ, de Jong M, Breit TM, Laine ML, de Soet JJ & Crielaard W (2010b) The core genome of the anaerobic oral pathogenic bacterium *Porphyromonas gingivalis*. *BMC Microbiol* **10**: 252.
- Carlisle MD, Srikantha RN & Brogden KA (2009) Degradation of human alpha- and beta-defensins by culture supernatants of *Porphyromonas gingivalis* strain 381. *J Innate Immun* **1**: 118–122.
- Chen T, Nakayama K, Belliveau L & Duncan MJ (2001) *Porphyromonas gingivalis* gingipains and adhesion to epithelial cells. *Infect Immun* **69**: 3048–3056.
- Chiang CY, Kyritsis G, Graves DT & Amar S (1999) Interleukin-1 and tumor necrosis factor activities partially account for calvarial bone resorption induced by local injection of lipopolysaccharide. *Infect Immun* **67**: 4231–4236.
- Choi JI, Schifferle RE, Yoshimura F & Kim BW (1998) Capsular polysaccharide-fimbrial protein conjugate vaccine protects against *Porphyromonas gingivalis* infection in SCID mice reconstituted with human peripheral blood lymphocytes. *Infect Immun* **66**: 391–393.
- Curtis MA, Thickett A, Slaney JM *et al.* (1999) Variable carbohydrate modifications to the catalytic chains of the RgpA and RgpB proteases of *Porphyromonas gingivalis* W50. *Infect Immun* **67**: 3816–3823.
- Curtis MA, Aduse-Opoku J & Rangarajan M (2001) Cysteine proteases of *Porphyromonas gingivalis*. *Crit Rev Oral Biol Med* **12**: 192–216.
- Darveau RP, Belton CM, Reife RA & Lamont RJ (1998) Local chemokine paralysis, a novel pathogenic mechanism for *Porphyromonas gingivalis*. *Infect Immun* **66**: 1660–1665.
- Darveau RP, Pham TT, Lemley K *et al.* (2004) *Porphyromonas gingivalis* lipopolysaccharide contains multiple lipid A species that functionally interact with both toll-like receptors 2 and 4. *Infect Immun* **72**: 5041–5051.
- Davey M, Liu X, Ukai T *et al.* (2008) Bacterial fimbriae stimulate proinflammatory activation in the endothelium through distinct TLRs. *J Immunol* **180**: 2187–2195.
- d'Empaire G, Baer MT & Gibson FC 3rd (2006) The K1 serotype capsular polysaccharide of *Porphyromonas gingivalis* elicits chemokine production from murine macrophages that facilitates cell migration. *Infect Immun* **74**: 6236–6243.
- Dierickx K, Pauwels M, Laine ML, Van Eldere J, Cassiman JJ, van Winkelhoff AJ, van Steenberghe D & Quirynen M (2003) Adhesion of *Porphyromonas gingivalis* serotypes to pocket epithelium. *J Periodontol* **74**: 844–848.
- Evans RT, Klausen B, Ramamurthy NS, Golub LM, Sfintescu C & Genco RJ (1992) Periodontopathic potential of two strains of *Porphyromonas gingivalis* in gnotobiotic rats. *Arch Oral Biol* **37**: 813–819.
- Fagundes JA, Monoo LD, Euzebio Alves VT, Pannuti CM, Cortelli SC, Cortelli JR & Holzhausen M (2011) *Porphyromonas gingivalis* is associated with protease-activated receptor-2 upregulation in chronic periodontitis. *J Periodontol* **82**: 1596–1601.
- Farquharson SI, Germaine GR & Gray GR (2000) Isolation and characterization of the cell-surface polysaccharides of *Porphyromonas gingivalis* ATCC 53978. *Oral Microbiol Immunol* **15**: 151–157.
- Fujise O, Hamachi T, Inoue K, Miura M & Maeda K (2002) Microbiological markers for prediction and assessment of treatment outcome following non-surgical periodontal therapy. *J Periodontol* **73**: 1253–1259.
- Gonzalez D, Tzianabos AO, Genco CA & Gibson FC 3rd (2003) Immunization with *Porphyromonas gingivalis* capsular polysaccharide prevents *P. gingivalis*-elicited oral bone loss in a murine model. *Infect Immun* **71**: 2283–2287.
- Grenier D, Roy E & Mayrand D (2003) Modulation of *Porphyromonas gingivalis* proteinase activity by suboptimal doses of antimicrobial agents. *J Periodontol* **74**: 1316–1319.
- Guo Y, Nguyen KA & Potempa J (2010) Dichotomy of gingipains action as virulence factors: from cleaving substrates with the precision of a surgeon's knife to a meat chopper-like brutal degradation of proteins. *Periodontology* **2000** **54**: 15–44.
- Haffajee AD, Cugini MA, Dibart S, Smith C, Kent RL Jr & Socransky SS (1997) The effect of SRP on the clinical and microbiological parameters of periodontal diseases. *J Clin Periodontol* **24**: 324–334.
- Hajishengallis G (2009) *Porphyromonas gingivalis*–host interactions: open war or intelligent guerilla tactics? *Microbes Infect* **11**: 637–645.
- Hajishengallis G & Lambris JD (2011) Microbial manipulation of receptor crosstalk in innate immunity. *Nat Rev Immunol* **11**: 187–200.
- Hajishengallis G, Tapping RI, Harokopakis E *et al.* (2006) Differential interactions of fimbriae and lipopolysaccharide from *Porphyromonas gingivalis* with the Toll-like receptor 2-centred pattern recognition apparatus. *Cell Microbiol* **8**: 1557–1570.
- Hajishengallis G, Shakhathreh MA, Wang M & Liang S (2007) Complement receptor 3 blockade promotes IL-12-mediated clearance of *Porphyromonas gingivalis* and negates its virulence *in vivo*. *J Immunol* **179**: 2359–2367.
- Hajishengallis G, Wang M, Liang S, Triantafyllou M & Triantafyllou K (2008) Pathogen induction of CXCR4/TLR2 cross-talk impairs host defense function. *P Natl Acad Sci USA* **105**: 13532–13537.
- Hajishengallis G, Wang M & Liang S (2009) Induction of distinct TLR2-mediated proinflammatory and proadhesive signaling pathways in response to *Porphyromonas gingivalis* fimbriae. *J Immunol* **182**: 6690–6696.
- Hajishengallis G, Liang S, Payne MA *et al.* (2011) Low-abundance biofilm species orchestrates inflammatory periodontal disease through the commensal microbiota and complement. *Cell Host Microbe* **10**: 497–506.
- Hamed M, Belibasakis GN, Cruchley AT, Rangarajan M, Curtis MA & Bostanci N (2009) *Porphyromonas gingivalis* culture supernatants differentially regulate interleukin-1beta and interleukin-18 in human monocytic cells. *Cytokine* **45**: 99–104.

- Holt SC, Kesavalu L, Walker S & Genco CA (1999) Virulence factors of *Porphyromonas gingivalis*. *Periodontology* **20**: 168–238.
- Holzhausen M, Cortelli JR, da Silva VA, Franco GC, Cortelli SC & Vergnolle N (2010) Protease-activated receptor-2 (PAR(2)) in human periodontitis. *J Dent Res* **89**: 948–953.
- Honda K (2011) *Porphyromonas gingivalis* sinks teeth into the oral microbiota and periodontal disease. *Cell Host Microbe* **10**: 423–425.
- Imamura T, Pike RN, Potempa J & Travis J (1994) Pathogenesis of periodontitis: a major arginine-specific cysteine proteinase from *Porphyromonas gingivalis* induces vascular permeability enhancement through activation of the kallikrein/kinin pathway. *J Clin Invest* **94**: 361–367.
- Imamura T, Potempa J, Pike RN & Travis J (1995a) Dependence of vascular permeability enhancement on cysteine proteinases in vesicles of *Porphyromonas gingivalis*. *Infect Immun* **63**: 1999–2003.
- Imamura T, Potempa J, Pike RN, Moore JN, Barton MH & Travis J (1995b) Effect of free and vesicle-bound cysteine proteinases of *Porphyromonas gingivalis* on plasma clot formation: implications for bleeding tendency at periodontitis sites. *Infect Immun* **63**: 4877–4882.
- Imamura T, Banbula A, Pereira PJ, Travis J & Potempa J (2001) Activation of human prothrombin by arginine-specific cysteine proteinases (Gingipains R) from *porphyromonas gingivalis*. *J Biol Chem* **276**: 18984–18991.
- Inaba H, Nakano K, Kato T *et al.* (2008) Heterogenic virulence and related factors among clinical isolates of *Porphyromonas gingivalis* with type II fimbriae. *Oral Microbiol Immunol* **23**: 29–35.
- Jagels MA, Ember JA, Travis J, Potempa J, Pike R & Hugli TE (1996a) Cleavage of the human C5A receptor by proteinases derived from *Porphyromonas gingivalis*: cleavage of leukocyte C5a receptor. *Adv Exp Med Biol* **389**: 155–164.
- Jagels MA, Travis J, Potempa J, Pike R & Hugli TE (1996b) Proteolytic inactivation of the leukocyte C5a receptor by proteinases derived from *Porphyromonas gingivalis*. *Infect Immun* **64**: 1984–1991.
- Jotwani R & Cutler CW (2004) Fimbriated *Porphyromonas gingivalis* is more efficient than fimbria-deficient *P. gingivalis* in entering human dendritic cells *in vitro* and induces an inflammatory Th1 effector response. *Infect Immun* **72**: 1725–1732.
- Jotwani R, Eswaran SV, Moonga S & Cutler CW (2010) MMP-9/TIMP-1 imbalance induced in human dendritic cells by *Porphyromonas gingivalis*. *FEMS Immunol Med Microbiol* **58**: 314–321.
- Kawada M, Yoshida A, Suzuki N, Nakano Y, Saito T, Oho T & Koga T (2004) Prevalence of *Porphyromonas gingivalis* in relation to periodontal status assessed by real-time PCR. *Oral Microbiol Immunol* **19**: 289–292.
- Kitamura Y, Matono S, Aida Y, Hirofujii T & Maeda K (2002) Gingipains in the culture supernatant of *Porphyromonas gingivalis* cleave CD4 and CD8 on human T cells. *J Periodontal Res* **37**: 464–468.
- Kolenbrander PE, Palmer RJ Jr, Periasamy S & Jakubovics NS (2011) Oral multispecies biofilm development and the key role of cell-cell distance. *Nat Rev Microbiol* **8**: 471–480.
- Kuboniwa M, Hasegawa Y, Mao S, Shizukuishi S, Amano A, Lamont RJ & Yilmaz O (2008) *P. gingivalis* accelerates gingival epithelial cell progression through the cell cycle. *Microbes Infect* **10**: 122–128.
- Kusumoto Y, Hirano H, Saitoh K *et al.* (2004) Human gingival epithelial cells produce chemotactic factors interleukin-8 and monocyte chemoattractant protein-1 after stimulation with *Porphyromonas gingivalis* via toll-like receptor 2. *J Periodontol* **75**: 370–379.
- Laine ML & van Winkelhoff AJ (1998) Virulence of six capsular serotypes of *Porphyromonas gingivalis* in a mouse model. *Oral Microbiol Immunol* **13**: 322–325.
- Laine ML, Appelmek BJ & van Winkelhoff AJ (1997) Prevalence and distribution of six capsular serotypes of *Porphyromonas gingivalis* in periodontitis patients. *J Dent Res* **76**: 1840–1844.
- Lamont RJ & Jenkinson HF (1998) Life below the gum line: pathogenic mechanisms of *Porphyromonas gingivalis*. *Microbiol Mol Biol Rev* **62**: 1244–1263.
- Liu X, Sroka A, Potempa J & Genco CA (2004) Coordinate expression of the *Porphyromonas gingivalis* lysine-specific gingipain proteinase, Kgp, arginine-specific gingipain proteinase, RgpA, and the heme/hemoglobin receptor, HmuR. *Biol Chem* **385**: 1049–1057.
- Liu R, Desta T, Raptis M, Darveau RP & Graves DT (2008) *P. gingivalis* and *E. coli* lipopolysaccharides exhibit different systemic but similar local induction of inflammatory markers. *J Periodontol* **79**: 1241–1247.
- Lourbakos A, Chinni C, Thompson P, Potempa J, Travis J, Mackie EJ & Pike RN (1998) Cleavage and activation of proteinase-activated receptor-2 on human neutrophils by gingipain-R from *Porphyromonas gingivalis*. *FEBS Lett* **435**: 45–48.
- Lourbakos A, Potempa J, Travis J *et al.* (2001) Arginine-specific protease from *Porphyromonas gingivalis* activates protease-activated receptors on human oral epithelial cells and induces interleukin-6 secretion. *Infect Immun* **69**: 5121–5130.
- Malek R, Fisher JG, Caleca A *et al.* (1994) Inactivation of the *Porphyromonas gingivalis* fimA gene blocks periodontal damage in gnotobiotic rats. *J Bacteriol* **176**: 1052–1059.
- McKee AS, McDermid AS, Baskerville A, Dowsett AB, Ellwood DC & Marsh PD (1986) Effect of hemin on the physiology and virulence of *Bacteroides gingivalis* W50. *Infect Immun* **52**: 349–355.
- Mydel P, Takahashi Y, Yumoto H *et al.* (2006) Roles of the host oxidative immune response and bacterial antioxidant rubrerythrin during *Porphyromonas gingivalis* infection. *PLoS Pathog* **2**: e76.
- Nakhjiri SF, Park Y, Yilmaz O, Chung WO, Watanabe K, El-Sabaeny A, Park K & Lamont RJ (2001) Inhibition of epithelial cell apoptosis by *Porphyromonas gingivalis*. *FEMS Microbiol Lett* **200**: 145–149.

- Nemoto E, Darveau RP, Foster BL, Nogueira-Filho GR & Somerman MJ (2006) Regulation of cementoblast function by *P. gingivalis* lipopolysaccharide via TLR2. *J Dent Res* **85**: 733–738.
- Nishida E, Hara Y, Kaneko T, Ikeda Y, Ukai T & Kato I (2001) Bone resorption and local interleukin-1 α and interleukin-1 β synthesis induced by *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* lipopolysaccharide. *J Periodontol Res* **36**: 1–8.
- Ogawa T, Ogo H, Uchida H & Hamada S (1994) Humoral and cellular immune responses to the fimbriae of *Porphyromonas gingivalis* and their synthetic peptides. *J Med Microbiol* **40**: 397–402.
- Oido-Mori M, Rezzonico R, Wang PL, Kowashi Y, Dayer JM, Baehni PC & Chizzolini C (2001) *Porphyromonas gingivalis* gingipain-R enhances interleukin-8 but decreases gamma interferon-inducible protein 10 production by human gingival fibroblasts in response to T-cell contact. *Infect Immun* **69**: 4493–4501.
- Paramonov N, Bailey D, Rangarajan M, Hashim A, Kelly G, Curtis MA & Hounsell EF (2001) Structural analysis of the polysaccharide from the lipopolysaccharide of *Porphyromonas gingivalis* strain W50. *Eur J Biochem* **268**: 4698–4707.
- Paramonov N, Rangarajan M, Hashim A *et al.* (2005) Structural analysis of a novel anionic polysaccharide from *Porphyromonas gingivalis* strain W50 related to Arg-gingipain glycans. *Mol Microbiol* **58**: 847–863.
- Paramonov NA, Aduse-Opoku J, Hashim A, Rangarajan M & Curtis MA (2009) Structural analysis of the core region of O-lipopolysaccharide of *Porphyromonas gingivalis* from mutants defective in O-antigen ligase and O-antigen polymerase. *J Bacteriol* **191**: 5272–5282.
- Paster BJ, Olsen I, Aas JA & Dewhirst FE (2006) The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontology 2000* **42**: 80–87.
- Pathirana RD, O'Brien-Simpson NM & Reynolds EC (2010) Host immune responses to *Porphyromonas gingivalis* antigens. *Periodontology 2000* **52**: 218–237.
- Pollreis A, Huang Y, Roth GA *et al.* (2010) Enhanced monocyte migration and pro-inflammatory cytokine production by *Porphyromonas gingivalis* infection. *J Periodontol Res* **45**: 239–245.
- Potempa J, Pike R & Travis J (1997) Titration and mapping of the active site of cysteine proteinases from *Porphyromonas gingivalis* (gingipains) using peptidyl chloromethanes. *Biol Chem* **378**: 223–230.
- Pulendran B, Kumar P, Cutler CW, Mohamadzadeh M, Van Dyke T & Banchereau J (2001) Lipopolysaccharides from distinct pathogens induce different classes of immune responses *in vivo*. *J Immunol* **167**: 5067–5076.
- Rangarajan M, Aduse-Opoku J, Paramonov N *et al.* (2008) Identification of a second lipopolysaccharide in *Porphyromonas gingivalis* W50. *J Bacteriol* **190**: 2920–2932.
- Sandros J, Karlsson C, Lappin DF, Madianos PN, Kinane DF & Papapanou PN (2000) Cytokine responses of oral epithelial cells to *Porphyromonas gingivalis* infection. *J Dent Res* **79**: 1808–1814.
- Schenkein HA, Fletcher HM, Bodnar M & Macrina FL (1995) Increased opsonization of a prtH-defective mutant of *Porphyromonas gingivalis* W83 is caused by reduced degradation of complement-derived opsonins. *J Immunol* **154**: 5331–5337.
- Schifferle RE, Reddy MS, Zambon JJ, Genco RJ & Levine MJ (1989) Characterization of a polysaccharide antigen from *Bacteroides gingivalis*. *J Immunol* **143**: 3035–3042.
- Scott CF, Whitaker EJ, Hammond BF & Colman RW (1993) Purification and characterization of a potent 70-kDa thiol lysyl-proteinase (Lys-gingivain) from *Porphyromonas gingivalis* that cleaves kininogens and fibrinogen. *J Biol Chem* **268**: 7935–7942.
- Sharma A, Honma K, Evans RT, Hruby DE & Genco RJ (2001) Oral immunization with recombinant *Streptococcus gordonii* expressing porphyromonas gingivalis FimA domains. *Infect Immun* **69**: 2928–2934.
- Shoji M, Ratnayake DB, Shi Y *et al.* (2002) Construction and characterization of a nonpigmented mutant of *Porphyromonas gingivalis*: cell surface polysaccharide as an anchorage for gingipains. *Microbiology* **148**: 1183–1191.
- Sims TJ, Schifferle RE, Ali RW, Skaug N & Page RC (2001) Immunoglobulin G response of periodontitis patients to *Porphyromonas gingivalis* capsular carbohydrate and lipopolysaccharide antigens. *Oral Microbiol Immunol* **16**: 193–201.
- Slaney JM, Gallagher A, Aduse-Opoku J, Pell K & Curtis MA (2006) Mechanisms of resistance of *Porphyromonas gingivalis* to killing by serum complement. *Infect Immun* **74**: 5352–5361.
- Smalley JW, Birss AJ, Szmigielski B & Potempa J (2006) The HA2 haemagglutinin domain of the lysine-specific gingipain (Kgp) of *Porphyromonas gingivalis* promotes micro-oxo bishaem formation from monomeric iron(III) protoporphyrin IX. *Microbiology* **152**: 1839–1845.
- Socransky SS (1979) Criteria for the infectious agents in dental caries and periodontal disease. *J Clin Periodontol* **6**: 16–21.
- Sundqvist G, Figdor D, Hanström L, Sorlin S & Sandström G (1991) Phagocytosis and virulence of different strains of *Porphyromonas gingivalis*. *Scand J Dent Res* **99**: 117–129.
- Takahashi Y, Davey M, Yumoto H, Gibson FC 3rd & Genco CA (2006) Fimbria-dependent activation of pro-inflammatory molecules in *Porphyromonas gingivalis* infected human aortic endothelial cells. *Cell Microbiol* **8**: 738–757.
- Tam V, O'Brien-Simpson NM, Chen YY, Sanderson CJ, Kinnear B & Reynolds EC (2009) The RgpA-Kgp proteinase-adhesin complexes of *Porphyromonas gingivalis* inactivate the Th2 cytokines interleukin-4 and interleukin-5. *Infect Immun* **77**: 1451–1458.
- Tribble GD, Mao S, James CE & Lamont RJ (2006) A *Porphyromonas gingivalis* haloacid dehalogenase family phosphatase interacts with human phosphoproteins and is important for invasion. *P Natl Acad Sci USA* **103**: 11027–11032.

- van Winkelhoff AJ, Loos BG, van der Reijden WA & van der Velden U (2002) *Porphyromonas gingivalis*, *Bacteroides forsythus* and other putative periodontal pathogens in subjects with and without periodontal destruction. *J Clin Periodontol* **29**: 1023–1028.
- Vernal R, Leon R, Silva A, van Winkelhoff AJ, Garcia-Sanz JA & Sanz M (2009) Differential cytokine expression by human dendritic cells in response to different *Porphyromonas gingivalis* capsular serotypes. *J Clin Periodontol* **36**: 823–829.
- Wang M, Shakhathreh MA, James D *et al.* (2007) Fimbrial proteins of *Porphyromonas gingivalis* mediate *in vivo* virulence and exploit TLR2 and complement receptor 3 to persist in macrophages. *J Immunol* **179**: 2349–2358.
- Wang M, Krauss JL, Domon H *et al.* (2010) Microbial hijacking of complement-toll-like receptor crosstalk. *Sci Signal* **3**: ra11.
- Wingrove JA, DiScipio RG, Chen Z, Potempa J, Travis J & Hugli TE (1992) Activation of complement components C3 and C5 by a cysteine proteinase (gingipain-1) from *Porphyromonas (Bacteroides) gingivalis*. *J Biol Chem* **267**: 18902–18907.
- Xia Q, Wang T, Taub F, Park Y, Capestany CA, Lamont RJ & Hackett M (2007) Quantitative proteomics of intracellular *Porphyromonas gingivalis*. *Proteomics* **7**: 4323–4337.
- Yang HW, Huang YF & Chou MY (2004) Occurrence of *Porphyromonas gingivalis* and *Tannerella forsythensis* in periodontally diseased and healthy subjects. *J Periodontol* **75**: 1077–1083.
- Yilmaz O (2008) The chronicles of *Porphyromonas gingivalis*: the microbium, the human oral epithelium and their interplay. *Microbiology* **154**: 2897–2903.
- Yilmaz O, Watanabe K & Lamont RJ (2002) Involvement of integrins in fimbriae-mediated binding and invasion by *Porphyromonas gingivalis*. *Cell Microbiol* **4**: 305–314.
- Yilmaz O, Young PA, Lamont RJ & Kenny GE (2003) Gingival epithelial cell signalling and cytoskeletal responses to *Porphyromonas gingivalis* invasion. *Microbiology* **149**: 2417–2426.
- Yilmaz O, Verbeke P, Lamont RJ & Ojcius DM (2006) Intercellular spreading of *Porphyromonas gingivalis* infection in primary gingival epithelial cells. *Infect Immun* **74**: 703–710.
- Yilmaz O, Yao L, Maeda K *et al.* (2008) ATP scavenging by the intracellular pathogen *Porphyromonas gingivalis* inhibits P2X7-mediated host-cell apoptosis. *Cell Microbiol* **10**: 863–875.
- Yun PL, DeCarlo AA & Hunter N (1999) Modulation of major histocompatibility complex protein expression by human gamma interferon mediated by cysteine proteinase-adhesin polyproteins of *Porphyromonas gingivalis*. *Infect Immun* **67**: 2986–2995.
- Yun PL, Decarlo AA, Collyer C & Hunter N (2001) Hydrolysis of interleukin-12 by *Porphyromonas gingivalis* major cysteine proteinases may affect local gamma interferon accumulation and the Th1 or Th2 T-cell phenotype in periodontitis. *Infect Immun* **69**: 5650–5660.
- Yun PL, DeCarlo AA, Collyer C & Hunter N (2002) Modulation of an interleukin-12 and gamma interferon synergistic feedback regulatory cycle of T-cell and monocyte cocultures by *Porphyromonas gingivalis* lipopolysaccharide in the absence or presence of cysteine proteinases. *Infect Immun* **70**: 5695–5705.
- Zhou Q, Desta T, Fenton M, Graves DT & Amar S (2005) Cytokine profiling of macrophages exposed to *Porphyromonas gingivalis*, its lipopolysaccharide, or its FimA protein. *Infect Immun* **73**: 935–943.
- Zijngel V, van Leeuwen MB, Degener JE, Abbas F, Thurnheer T, Gmur R & Harmsen HJ (2011) Oral biofilm architecture on natural teeth. *PLoS ONE* **5**: e9321.